## THE USE OF OUTCOME MEASURES WITH CHILDREN/YOUNG PEOPLE WITH MORE SEVERE LEARNING DISABILITIES: What works in practice?

### UPDATE from Workshop #1 (October 2014)

#### **Alex Crawford/Neil Phillips**

Worcestershire CAMHS Learning Disability Team Worcestershire Health and Care NHS Trust

#### Sara Sopena

Wandsworth CAMHS Learning Disability Team South West London and St Georges NHS Trust


12<sup>th</sup> March 2015


## UPDATE


# Preliminary results


- Demographics
  - N = 94
  - 51 clinicians gave their details
  - 30 said they would be interested to be part of a bigger study

### Profession (N=51)









### Do you use this measure with families with children with LD?



## Specific answers

	Useful clinically? %	Parents find it useful? %	Easy to complete? %
Goals-based	97	92	88
SLDOM	88	71	82
CHI-ESQ	93	88	93
SDQ	70	37	60
DBC	90	59	54

# Other measures reported

Name of outcome measure	Reported by no of services
Aberrant behaviour Checklist	1
Behaviour grids to measure parental perceptions of difficulties	2
Checklist of Challenging Behaviour and Rating Scales	1
Complex Sleep disturbance index	1
Conners	1
Considering a skills questionnaire such as "Essentials for Living"	1
Current View	1
HADS for parental mental health	1
Honosca	2
Paddington Complexity Scale.	1
Parenting competence scale	1
Parents top 3 concerns	3
PedsQOL battery	1
Questionnaire on Resources and Stress.	2
RCADS for IAPT	3
Service developed satisfaction questionnaire	1
Sleep specific measures	1
Social communication questionnaire	1

# Conclusions

On the basis of this survey the suggested outcome measures to use are:

- Focus of work:
  - Goals-based outcome measure
- Parenting competence/understanding/confidence:
  - SLDOM
- Behavioural/emotional needs of child:
  - DBC
- Satisfaction:
  - CHI-ESQ

# Conclusions

- However, need to:
  - Evaluate effectiveness of OM more systematically (pre and post intervention?, service user feedback?).
  - In line with IAPT principles.
  - Look at other measures that are not commonly used but have been rated as being useful.
  - Need to develop new measures?

# Next Steps

### • Evaluate effectiveness of OM more systematically:

- Services to volunteer to evaluate OM pre and post intervention
- Get service user feedback (as part of same study?)
- In line with IAPT principles.
  - To discuss today?
- Look at other measures that are not commonly used but have been rated as being useful.
  - Subsequent project if needed?
- Need to develop new measures?
  - Subsequent project if needed?

## Contact details

Neil Phillips

neil.phillips@hacw.nhs.uk 01684 612740

• Sara Sopena

sara.sopena@nhs.net

077 532 05 237